

Phosphorus Forms at Different Spatial Scales in the Lower Fox River Sub-Basin

Nicholas A. Reckinger, UW-Green Bay, reckna20@uwgb.edu

Kevin Fermanich, UW-Green Bay

Paul Baumgart, UW-Green Bay

In many settings, particulate phosphorus (PP) has been reported to be the dominant form of P in surface water runoff from agricultural systems. Past monitoring of rural streams in the Lower Fox River sub-basin in Northeastern Wisconsin has shown mean concentrations of dissolved phosphorus representing from 40 to 75% of total P (TP) concentration. The effectiveness of improved management strategies that target PP is dependent on the form of P leaving source areas. Total dissolved phosphorus (TDP) losses may not be reduced with best management practices such as grassed water ways, filter strips, and conservation tillage. This study was conducted to better understand P forms leaving source areas and how the dissolved P fraction changes along a flow path at different scales in the Apple Creek watershed, Outagamie County, Wisconsin. Samples were collected near peak flow at 11 rural source area sites (0.25 to 2.5 km²) and 4 integrator sites (12 to 85 km²) during runoff events in 2004-06. A USGS continuous monitoring station also collected event samples on the main stem (117 km²). For six runoff events in 2004-05, median TP was 0.51 mg/L from source area sites, 0.48 mg/L from integrator sites, and 0.72 mg/L from the main stem. Median TDP percentage was 39% from source areas, 40% from integrator sites, and 27% at the main stem. The median TDP percentage for the six events at each source area site, varied greatly (10% to 82%). The portion of TDP in a snowmelt and a low intensity event in 2006 were twice the median from earlier events.